DIRECTIONS FOR USE

Boom spray, Mist spray or Aerial

Use appropriate personal protective equipment and handling equipment.

Agitate contents well before use.

Fill tank to a minimum of two thirds of water.

Add BioAg Balance & Grow under agitation, bring volume up to full. Before tank mixing, jar test all added nutrients for compatibility of use. Check for physical changes.

Add other nutrients as prescribed under agitation.

Use within 48 hours of dilution.

Rinse tanks and sprayer and flush lines after use.

TYPICAL RATES OF APPLICATION

Crop	Rate L/ha	Notes	
Broad Acre Grasses Cereals, sorghum, corn, sugar cane	1.0 to 3.0	Apply to non-stressed plants from early tillering until first internode elongation of 25 mm. Do not apply after booting commences.	
Broad Acre Dicots Cotton, potatoes, clovers, lucerne, peanuts, canola, pulses, sunflowers	1.0 to 3.5 1 to 4 Applications	Apply to non-stressed plants during the vegetative growth stage.	
Horticulture Vines, citrus, vegetables, nut and fruit trees	1.0 to 4.0 1 to 6 Applications	Apply to plants prior to the commencement of their reproductive phase at a minimum dilution of 1:100 with water.	
Pastures	2.0 to 4.0	Apply to grazed pastures. Withholding period – NIL	

NOTE: The above rates should be used as a guide only. BioAg does not warrant that these rates will necessarily produce the optimal or expected outcome. Each farmer's climatic conditions, before, during and after application, water quality, soil and plant types, application processes and practices may differ and require adjustments to the application rates to ensure optimum results.

Manufactured by BioAg Pty Ltd ABN 58 086 880 211 22-26 Twynam St Narrandera NSW 2700 Australia T: +61 2 6958 9911 F: +61 2 6959 9922 W: www.bioag.com.au

KEEP OUT OF REACH OF CHILDREN



Balance & Grow[®] Organic[#]

Foliar Growth Stimulant







Australian made, owned and operated # allowed for use in organic production systems

ABOUT BALANCE & GROW

Produced by aerobic fermentation, BioAg Balance & Grow has been developed to increase the vegetative growth of treated plants and to aid in the expansion of the leaf and soil foodweb. Contact BioAg for a product analysis.

WARNING STATEMENTS

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

STORAGE AND HANDLING

Store out of direct sunlight and below 30°C, away from children, animals, food and feed. Do not store mixed product. Shelf life of 36 months when stored per directions.

CONDITIONS OF SALE:

The product has a limited shelf life and must be stored in accordance with the storage and handling directions to preserve shelf life. Failure to store correctly or use after the expiry of shelf life is a breach of the conditions of sale. As BioAg cannot exercise control over storage, handling, and the exercise of personal judgment, skill and expertise in the mixing, application or use, or weather, or animal condition before, during or after application, all of which may affect the performance of the product, BioAg, to the maximum extent permitted by law, will not accept any liability for any failure in performance of the product, or any losses or damages whatsoever.

Except as expressly provided in these Conditions of Sale, no warranty, condition, undertaking or term, express or implied, statutory or otherwise, as to the quality or fitness for purpose of the goods is given or assumed by BioAg and all such conditions, warranties and liabilities (including liability as to negligence) express or implied relating to the goods are hereby excluded to the extent permitted by law. To the extent that liability can be shown (under law) it is limited to the replacement of goods or their equivalent monetary value only.

Batch No / DOM:					
Net Contents	20 L	200 L	1,000 L		