

Pasture Primo™

0-3.8-0-0

Biologically treated phosphate with lime



FOR SOILS THAT REQUIRE PHOSPHORUS AND
CALCIUM.

Users are encouraged to seek further advice:
www.bioag.com.au/agronomy-sales.

Read label before use.

TYPICAL ANALYSIS (DRY WEIGHT)

Total Phosphorus (P)	3.8%
Phosphorus – Citric Soluble (2Hr)#	1.4%
Phosphorus – Formic Soluble (2Hr)*	2.7%
Calcium as phosphate/carbonate/oxide (Ca)	37%
Sulphur as sulphate	0.3%
Cadmium [Maximum]	4.9 ppm (131 mg/kg P)
Cadmium [Maximum Allowed Limit]	300 mg/kg P
Fluorine	1.1% (281 g/kg P)

SIEVE SPECIFICATIONS

>1 mm	2%
-------	----

PHYSICAL DESCRIPTION

Light grey to off white sand.

APPLICATION

Applied using a belt spreader. Risk of blow under windy conditions.

APPLICATION RATE

Consult with your BioAg agronomist or distributor.

* Citric Acid and Formic Acid tests performed in accordance with methods used in the CSIRO 1997 National Reactive Phosphate Rock Project

SAFETY DIRECTIONS

Do not swallow. The dust from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Wear protective gloves, protective clothing and eye/face protection. Wash contaminated skin after handling.

STORAGE & HANDLING

Store dry conditions.

COMPATIBILITY

Can be mixed with gypsum.

FIRST AID

For advice, contact a Poisons Information Centre (e.g. phone Australia 13 11 26; New Zealand 0800 764 766) or a doctor (at once). If in eyes flush with plenty of water for 10 - 15 minutes. Keep eyes wide open while flushing. Seek medical attention if irritation persists.

ADDITIONAL INFORMATION

Dispose of contents & container in accordance with local regulations. Refer to your BioAg Pty Ltd advisor, the BioAg website (www.bioag.com.au) or the SDS.

Manufactured by BioAg Pty
Ltd 22-26 Twynam St
Narrandera NSW 2700
T: +61 2 6958 9911
W: www.bioag.com.au



Better soils. Better crops. Better stock.™